



In April of 2008 a panel of internationally recognized experts confirmed that Reb A is safe to use as a table top sweetener and as an ingredient in beverages and cereals. The panel also concluded that Reb A meets the Generally Recognized as Safe (GRAS) criteria established by the Food and Drug Administration (FDA). Science supporting the safety of Reb A includes:

1. Abudula, R., Jeppesen, P.B., Rolfsen, S.E., Xiao, J., and Hermansen, K. 2004. Rebaudioside A potently stimulates insulin secretion from isolated mouse islets: studies on the dose-, glucose-, and calcium-dependency. *Metabolism*. 53(10):1378-1381.
2. Aze, Y., Toyoda, K., Imaida, K., Hayashi, S., Imazawa, T., Hayashi, Y. and Takahashi, M. 1991. Subchronic Oral Toxicity Study of Stevioside in F344 Rats. *Eisei Shikenjo Hokoku (Japanese)*. 3(109):48-54.
3. Barriocanal, L.A., Palacios, M., Benitez, G., Benitez, S., Jimenez, J.T., Jimenez, N., and Rojas, V. 2008. Apparent Lack of Pharmacological Effect of Steviol Glycosides Used as Sweeteners in Humans. A Pilot Study of Repeated Exposures in Some Normotensive and Hypotensive Individuals and in Type 1 and Type 2 Diabetics. *Regul Toxicol Pharmacol*. In Press, Available online 5 March 2008.
4. Brusick, D.J. 2008. A critical review of the genetic toxicity of steviol and steviol glycosides. *Food Chem Toxicol*. 46(7)Suppl: S83-S91.
5. Chan, P., Tomlinson, B., Chen, Y.J., Liu, J.C., Hsieh, M.H., and Cheng, J.T. 2000. A double-blind placebo-controlled study of the effectiveness and tolerability of oral stevioside in human hypertension. *Br J Clin Pharmacol*. 50:215-220.
6. Curi, R., Alvarez, M., Bazotte, R.B., Botion, L.M., Godoy, J.L., and Bracht, A. 1986. Effect of *Stevia rebaudiana* on glucose tolerance in normal adult humans. *Braz J Med Biol Res*. 19:771-4.
7. Curry, L.L., Roberts, A. 2008. Subchronic toxicity of rebaudioside A. *Food Chem Toxicol*. 46(7)Suppl: S11-S20.
8. Curry, L.L., Roberts, A., Brown, N. 2008. Rebaudioside A: two-generation reproductive toxicity study in rats. *Food Chem Toxicol*. 46(7)Suppl: S21-S30.
9. Dyrskog, S.E.U., Jeppesen, P.B., Chen, J., Christensen, L.P., and Hermansen, K. 2005a. The diterpene glycoside, Rebaudioside A, does not improve glycemic control or affect blood pressure after eight weeks treatment in the Goto-Kakizaki rat. *Rev Diabet Stud*. 2: 84-91.

10. Dyrskog, S.E.U., Jeppesen, P.B., Colombo, M. Abudula, R. and Hermansen, K. 2005b. Preventive effects of a soy-based diet supplemented with stevioside on the development of the metabolic syndrome and type 2 diabetes in Zucker diabetic fatty rats. *Metabolism*. 54:1181-1188.
11. Ferri, L.A.F., Alves-Do-Prado, W., Yamada, S.S., Gazola, S., Batista, M.R., and Bazotte, R.B. 2006. Investigation of the antihypertensive effect of oral crude stevioside in patients with mild essential hypertension. *Phytother Res*. 20:732-736.
12. Flamm, W.G., Blackburn, G.L., Comer, C.P., Mayhew, D.A. and Stargel, W.W. 2003. Long-term food consumption and body weight changes in neotame safety studies are consistent with the allometric relationship observed for other sweeteners and during dietary restrictions. *Regul Toxicol Pharmacol*. 38:144-156.
13. FSANZ (Food Standards Australia New Zealand). 2007. Steviol Glycosides as Intense Sweeteners. Draft Assessment Report Application A540, pp. 1-89.
14. Gardana, C., Simonetti, P., Canzi, E., Zanchi, R., and Pietta, P. 2003. Metabolism of stevioside and rebaudioside A from *Stevia rebaudiana* extracts by human microflora. *J Agric Food Chem*. 51:6618-22.
15. Geuns, J.M.C. 2007. Letter to the Editor on: Nunes, A.P.M., Ferreira-Machado, S.C., Nunes, R.M., Dantas, F.J.S., DeMattos, J.C.P.D., Caldeira-de-Araujo, A., 2007. Analysis of genotoxic potentiality of stevioside by comet assay. *Food Chem Toxicol*. 45: 662–666, 2597–2598, 2601–2602.
16. Geuns, J.M.C. and Pietta, P. 2004. Stevioside Metabolism by Human Volunteers. From symposium.
17. Geuns, J.M.C., Augustijns, P., Mols, R., Buyse, J.G., and Driessen, B. 2003a. Metabolism of stevioside in pigs and intestinal absorption characteristics of stevioside, rebaudioside A and steviol. *Food Chem Toxicol*. 41:1599-607.
18. Geuns, J.M.C., Malheiros, R.D., Moraes, V.M., Decuypere, E.M., Compennolle, F. and Buyse, J.G. 2003b. Metabolism of stevioside by chickens. *J Agric Food Chem*. 51:1095-101.
19. Geuns, J.M.C., Buyse, J., Vankeirsbilck, A., Temme, E.H.M. 2006. Compennolle, F. and Toppet, S. Identification of steviol glucuronide in human urine. *J Agric Food Chem*. 54:2794–2798.
20. Geuns, J.M.C., Buyse, J., Vankeirsbilck, A., and Temme, E.H.M. 2007. Metabolism of stevioside by healthy subjects. *Exp Biol Med*. 232:164-173.
21. GRAS PANEL. 2008. GRAS PANEL OPINION on the GRAS status of Reb A. GRN No. 252. (<http://www.cfsan.fda.gov/~rdb/opa-gn08.html>).
22. Gregersen, S., Jeppesen, P.B., Holst, J.J., and Hermansen, K. 2004. Antihyperglycemic effects of stevioside in type 2 diabetic subjects. *Metabolism*. 53:73-76.

23. Haebisch, B. 1992. Pharmacological trial of a concentrated crude extract of stevia rebaudiana (bert.) bertonii in healthy volunteers. *Arq Biol Technol.* 35:299-314.
24. Hanson, J.R. and De Oliveira, B.H. 1993. Stevioside and related sweet diterpenoid glycosides. *Nat Prod Rep.* 10(3):301- 309.
25. Hsieh, M.H., Chan, P., Sue, Y.M., Liu, J.C., Liang, T.H., Huang, T.Y., Tomlinson, B., Chow, M.S., Kao, P.F., and Chen, Y.J. 2003. Efficacy and tolerability of oral stevioside in patients with mild essential hypertension: a two-year, randomized, placebo-controlled study. *Clin Ther.* 25:2797-2808.
26. Hutapea, A.M., Toskulkao, C., Buddhasukh, D., Wilairat, P., Glinsukon, T. 1997. Digestion of stevioside, a natural sweetener, by various digestive enzymes. *J Clin Biochem Nutr.* 23:177–186.
27. JECFA (Joint FAO/WHO Expert Committee on Food Additives). 1999. Safety evaluation of certain food additives. WHO Food Additives Series 42. World Health Organization, Geneva. pp. 119-143.
28. JECFA (Joint FAO/WHO Expert Committee on Food Additives). 2006. Compendium of food additive specifications. Steviol glycosides. New tentative specifications prepared at the Sixty-third Report of the Joint FAO/WHO Expert Committee on Food Additives.
29. JECFA (Joint FAO/WHO Expert Committee on Food Additives). 2007. Compendium of food additive specifications. World Health Organization, Geneva. Steviol glycosides. Specifications prepared at the Sixty-eighth Report of the Joint FAO/WHO Expert Committee on Food Additives.
30. JECFA (Joint FAO/WHO Expert Committee on Food Additives). 2008. Sixty-ninth meeting Rome, Italy, 17-26 June 2008. Summary and conclusions.
31. Jeppesen, P.B., Gregersen, S., Poulsen, C.R., and Hermansen, K. 2000. Stevioside acts directly on pancreatic beta cells to secrete insulin: actions independent of cyclic adenosine monophosphate and adenosine triphosphate-sensitive K⁺-channel activity. *Metabolism.* 49:208-214.
32. Jeppesen, P.B., Gregersen, S., Alstrup, K.K., and Hermansen, K. 2002. Stevioside induces antihyperglycaemic, insulinotropic and glucagonostatic effects *in vivo*: studies in the diabetic Goto-Kakizaki (GK) rats. *Phytomedicine.* 9:9-14.
33. Jeppesen, P.B., Gregersen, S., Rolfsen, S.E., Jepsen, M., Colombo, M., Agger, A., Xiao, J., Kruhoffer, M., Orntoft, T., and Hermansen, K. 2003. Antihyperglycemic and blood pressure-reducing effects of stevioside in the diabetic Goto-Kakizaki rat. *Metabolism.* 52:372-378.
34. Jeppesen, P.B., Barriocanal, L., Meyer, M.T., Palacios, M., Canete, F., Benitez, S., Logwin, S., Schupmann, Y., Benitez, G., and Jimenez, J.T. 2006. Efficacy and tolerability of oral stevioside in patients with type 2 diabetes: a long-term, randomized, double-blinded, placebo-controlled study. *Diabetologia.* 49:1-755.
35. JFJA (Japan Fruit Juice Association). 2007. *The Unified Test Method for Thermo-Acidophilic Bacilli.* Japan Fruit Juice Association, Tokyo, Japan.

36. Klongpanichpak, S., Temcharoen, P., Toskulkao, C., Apibal, S., and Glinsukon, T. 1997. Lack of mutagenicity of stevioside and steviol in *Salmonella typhimurium* TA98 and TA100. *J Med Assoc Thai.* 80(Suppl 1):S121-188.
37. Koyama, E., Sakai, N., Ohori, Y., Kitazawa, K., Izawa, O., Kakegawa, K., Fujino, A., and Ui, M. 2003a. Absorption and metabolism of the glycosidic sweeteners of *Stevia* mixture and their aglycone, steviol, in rats and humans. *Food Chem Toxicol.* 41:875–883.
38. Koyama, E., Kitazawa, K., Ohori, Y., Izawa, O., Kakegawa, K., Fujino, A., Ui, M. 2003b. In vitro metabolism of the glycosidic sweeteners, *Stevia* mixture and enzymatically modified *Stevia* in human intestinal microflora. *Food Chem Toxicol.* 41:359–374.
39. Liu, J.C., Kao, P.F., Hsieh, M.H., Chen, Y.J., and Chan, P. 2001. The antihypertensive effects of stevioside derivative Isosteviol in spontaneously hypertensive rats. *Acta Cardiologica Sinica.* 17:133–140. As cited in Xu *et al.* (2007b).
40. Liu, J.C., Kao, P.K., Chan, P., Hsu, Y.H., Hou, C.C., Lien, G.S., Hsieh, M.H., Chen, Y.J., and Cheng, J.T. 2003. Mechanism of the antihypertensive effect of stevioside in anesthetized dogs. *Pharmacology.* 67:14-20.
41. Maki, K.C., Curry, L.L., Carakostas, M.C., Tarka, S.M., Reeves, M.S., Farmer, M.V., McKenney, J.M., Toth, P.D., Schwartz, S.L., Lubin, B.C., Dicklin, M.R., Boileau, A.C., Bisognano, J.D. 2008a. The hemodynamic effects of rebaudioside A in healthy adults with normal and low-normal blood pressure. *Food Chem Toxicol.* 46(7)Suppl: S40-S46.
42. Maki, K.C., Curry, L.L., Reeves, M.S., Toth, P.D., McKenney, J.M., Farmer, M.V., Schwartz, S.L., Lubin, B.C., Boileau, A.C., Dicklin, M.R., Carakostas, M.C., Tarka, S.M. 2008b. Chronic consumption of rebaudioside A, a steviol glycoside, in men and women with type 2 diabetes mellitus. *Food Chem Toxicol.* 46(7)Suppl: S47-S53.
43. Matsui, M., Matsui, K., Nohmi, T., Mizusawa, H., and Ishidate, M., Jr. 1989. Detection of deletion mutations in pSV@gpt plasmids induced by metabolically activated steviol. *Mutat Res.* 216:367-368.
44. Matsui, M., Matsui, K., Kawasaki, Y., Oda, Y., Noguchi, T., Kitagawa, Y., Sawada, M., Hayashi, M., Nohmi, T., Yoshihira, K., Ishidate, M., Jr., and Sofuni, T. 1996a. Evaluation of the genotoxicity of stevioside and steviol using six *in vitro* and one *in vivo* mutagenicity assays. *Mutagenesis.* 11:573-579.
45. Matsui, M., Sofuni, T., and Nohmi, T. 1996b. Regionally-targeted mutagenesis by metabolically-activated steviol: DNA sequence analysis of steviol-induced mutants of guanine phosphoribosyltransferase (gpt) gene of *Salmonella typhimurium* TM677. *Mutagenesis.* 11:565-572.
46. Medon, P., Pezzuto, J.M., Hovanec-Brown, J., Nanayakkara, D., Soejarto, D., Kamath, S., and Kinghorn, A.D. 1982. Safety assessment of some *Stevia rebaudiana* sweet principles. Fed. Proc. 41:1568 (Abstract). Federation of American Societies for Experimental Biology: 66th Annual Meeting, New Orleans, LA.

47. Melis, M.S. 1995. Chronic administration of aqueous extract of *Stevia rebaudiana* in rats: renal effects. *J Ethnopharmacol.* 47:129-134.
48. Melis, M.S. 1996. A crude extract of *Stevia rebaudiana* increases the renal plasma flow of normal and hypertensive rats. *Braz J Med Biol Res.* 29:669-675.
49. Melis, M.S. 1999. Effects of chronic administration of *Stevia rebaudiana* on fertility in rats. *J Ethnopharmacol.* 67:157-161.
50. Nakayama, K., Kasahara, D., and Yamamoto, F. 1986. Absorption, distribution, metabolism and excretion of stevioside in rats. *J Food Hyg Soc Jpn.* 27:1-8.
51. NCHS (National Center for Health Statistics). 2007. Centers for Disease Control and Prevention (CDC). National Health and Nutrition Examination Survey Data 2003-2004. Hyattsville, MD: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, 2007 (http://www.cdc.gov/nchs/about/major/nhanes/nhanes2003-2004/nhanes03_04.htm).
52. Nikiforov, A.I. and Eapen, A.K. 2008. A 90-day oral (dietary) toxicity study of rebaudioside A in Sprague-Dawley rats. *Int J Toxicol.* 27(1):65-80.
53. Oh, H.-Y., Han, E.-S., Choi, D.-W., Kim, J.-W., Eom, M.-O., Kang, I.-H., Kang, H.-J., and Ha, K.-W. 1999. *In vitro* and *in vivo* evaluation of genotoxicity of stevioside and steviol, natural sweetener. *Yakhak Hoeji.* 43(5):614-622.
54. Pezzuto, J.M., Compadre, C.M., Swanson, S.M., Nanayakkara, N.P.D., and Kinghorn, A.D. 1985a. Metabolically activated steviol, the aglycone of stevioside, is mutagenic. *Proc Natl Acad Sci USA.* 82:2478-2482.
55. Pezzuto, J.M., Nanayakkara, N.P.D., Compadre, C.M., Swanson, S.M., Kinghorn, A.D., Guenther, T.M., Sparnins, V.L., and Lam, L.K. 1986. Characterization of bacterial mutagenicity mediated by 13-hydroxy-ent-kaurenoic acid (steviol) and several structurally-related derivatives and evaluation of potential to induce glutathione S-transferase in mice. *Mutat Res.* 169:93-103.
56. Procinska, E., Bridges, B.A., and Hanson, J.R. 1991. Interpretation of results with the 8-azaguanine resistance system in *Salmonella typhimurium*: no evidence for direct acting mutagenesis by 15-oxosteviol, a possible metabolite of steviol. *Mutagenesis.* 6:165-167.
57. Renwick, A.G. 2008. The use of a sweetener substitution method to predict dietary exposures for the intense sweetener rebaudioside A. *Food Chem Toxicol.* 46(7)Suppl: S61-S69.
58. Roberts, A., Renwick, A.G. 2008. Comparative toxicokinetics and metabolism of rebaudioside A, stevioside, and steviol in rats. *Food Chem Toxicol.* 46(7)Suppl: S31-S39.
59. Simonetti, P., Gardana, C., Bramati, L., and Pietta, P.G. 2004. Bioavailability of Stevioside from *Stevia rebaudiana* in human volunteers: preliminary report. In: Geuns J.M.C. and Buyse, J., Eds. Proceedings of the first symposium on the safety of stevioside, KULeuven, April 16, 2004. Heverlee-Leuven, Belgium: Euprint Editions, pp. 51-62.

60. Suanarunsawat, T. and Chaiyabutr, N. 1997. The effect of stevioside on glucose metabolism in rat. *Can J Physiol Pharmacol.* 75:976-982.
61. Suttajit, M., Vinitketkaumnue, U., Meevatee, U., and Buddhasukh, D. 1993. Mutagenicity and human chromosomal effect of stevioside, a sweetener from *Stevia rebaudiana* Bertoni. *Environ Health Perspect.* 101(Suppl 3):53-56.
62. Takanaka, T., Kawashima, K., Usami, M. and Sakami, K. 1991. A teratological study of stevioside administered orally to rats. Unpublished report from Department of Pharmacology, Biological Safety Research Center, National Institute of Hygienic Sciences, Japan. Submitted to WHO by Ministry of Health and Welfare, Food Chemistry Division, Japan. As cited in JECFA (1999).
63. Temcharoen, P., Suwannatrai, M., Klongpanichpak, S., Apibal, S., Glinsukon, T., and Toskulkao, C. 2000. Evaluation of the effect of steviol on chromosomal damage using micronucleus test in three laboratory animal species. *J Med Assoc Thai.* 83(Suppl 1):S101-8.
64. Terai, T., Ren, H., Mori, G., Yamaguchi, Y., and Hayashi, T. 2002. Mutagenicity of steviol and its oxidative derivatives in *Salmonella typhimurium* TM677. *Chem Pharm Bull (Tokyo).* 50:1007-1010.
65. Toskulkao, C., Chaturat, L., Temcharoen, P., and Glinsukon, T. 1997. Acute toxicity of stevioside, a natural sweetener, and its metabolite, steviol, in several animal species. *Drug Chem Toxicol.* 20:31-44.
66. Toyoda, K., Matsui, H., Shoda, T., Uneyama, C., Takada, K., and Takahashi, M. 1997. Assessment of the carcinogenicity of stevioside in F344 rats. *Food Chem Toxicol.* 35:597-603.
67. Usami, M., Sakemi, K., Kawashima, K., Tsuda, M., and Ohno, Y. 1995. Teratogenicity study of stevioside in rats. *Bull Natl Inst Hyg Sci.* 113:31-35.
68. Wasuntarawat, C., Temcharoen, P., Toskulkao, C., Mungkornkarn, P., Suttajit, M., and Glinsukon, T. 1998. Developmental toxicity of steviol, a metabolite of stevioside, in the hamster. *Drug Chem Toxicol.* 21:207-22.
69. Wheeler, A., Boileau, A.C., Winkler, P.C., Compton, J.C., Prakash, I., Jiang, X., and Mandarino, D.A. 2008. Pharmacokinetics of rebaudioside A and stevioside after single oral doses in healthy men. *Food Chem Toxicol.* 46(7)Suppl: S54-S60.
70. Wingard, R.E., Jr., Brown, J.P., Enderlin, F.E., Dale, J.A., Hale, R.L., and Seitz, C.T. 1980. Intestinal degradation and absorption of the glycosidic sweeteners stevioside and rebaudioside A. *Experientia.* 36:519-520.
71. Xili, L., Chengjian, B., Eryi, X., Reiming, S., Yuengming, W., Haodong, S., and Zhiyian, H. 1992. Chronic oral toxicity and carcinogenicity study of stevioside in rats. *Food Chem Toxicol.* 30:957-965.